

Seton Hall University

eRepository @ Seton Hall

Petersheim Academic Exposition

Petersheim Academic Exposition

4-2020

Adjacency Eigenvalues for Underlying Split Multigraphs

Analisa Espino

Seton Hall University

Follow this and additional works at: <https://scholarship.shu.edu/petersheim-exposition>



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Espino, Analisa, "Adjacency Eigenvalues for Underlying Split Multigraphs" (2020). *Petersheim Academic Exposition*. 91.

<https://scholarship.shu.edu/petersheim-exposition/91>



Adjacency Eigenvalues for Underlying Split Multigraphs

ANALISA ESPINO

SETON HALL UNIVERSITY, UNDERGRADUATE STUDENT
DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE


MENTOR: DR. SACCOMAN

Background Information

$$x = IPS(c, d)^\mu$$

- c = number of cones
- d = degree of each cone
- μ = multiplicity of edges within clique
- x = number of cone nodes to which each clique node is adjacent

$$2 = IPS(8, 2)^\mu$$



Research Question: If these multigraphs represent satellite and ground station communication, can we find a formula to best represent the number of triangles from the ground station to the satellite?

Method

Characteristic Polynomial

- Submatrix:

```
CharacteristicPolynomial[CP91, x]
```

$$8 + 63x + 216x^2 + 420x^3 + 504x^4 + 378x^5 + 168x^6 + 36x^7 - x^9$$

168 / 2

84

- Whole Matrix:

```
CharacteristicPolynomial[IPS931, x]
```


$$-729x^6 - 4860x^7 - 13473x^8 - 19502x^9 - 14427x^{10} - 2862x^{11} + 3360x^{12} + 2358x^{13} + 243x^{14} - 222x^{15} - 63x^{16} + x^{18}$$

222 / 2

111

111 - 84

27



Conjecture


1. Number of triangles can be found with the formula: $\frac{(x-1)}{2} * (x^2 * c_0 * \mu)$
2. $c_0(x - 1) =$ number of eigenvalues equalling zero
3. $c_0(x - 1) =$ number of eigenvalues equalling $-\mu$
4. There are two eigenvalues that are the roots of $x^2 + \mu x + -(x^2)$
5. There is one eigenvalue that is the roots of $x^2 + (-\mu)(c - 1)x - x^2$

Triangles


- Conjecture 1: $\frac{(x-1)}{2} * (x^2 * c_o * \mu)$

$$\frac{(3-1)}{2} * (3^2 * 3 * 1)$$
$$1 * (9 * 1)$$
$$27$$

* Note: $c = x * c_o$
 $9 = 3 * 3$



Eigenvalues



{9., -3.54138, -3.54138, 2.54138, 2.54138, -1., -1., -1., -1., -1., -1., -1., 0., 0., 0., 0., 0., 0.}

Eigenvalues = 0 have a multiplicity of 6
 Eigenvalues = -1 have a multiplicity of 6


$$3 - IPS(9, 3)^1$$

$$c_o = 3$$

$$x = 3$$

Conjecture 2 & 3: $c_o(x - 1) = \text{number of eigenvalues equalling zero \& equalling } -\mu$

Eigenvalues



- Conjecture 4:
There are two eigenvalues that are the roots of $x^2 + \mu x + -(x^2)$

{9., -3.54138, -3.54138, 2.54138, 2.54138, -1., -1., -1., -1., -1., -1., -1., 0., 0., 0., 0., 0., 0.}

$$\begin{aligned} x^2 + \mu x + (-x^2) \\ 1x^2 + 1x - 9 \end{aligned}$$


Factored Polynomial Quotient: $(x^2 + x - 9)$

$$\begin{aligned} A &= 1 \\ B &= 1 \\ C &= -9 \end{aligned}$$

$$\frac{(B \pm \sqrt{B^2 - 4AC})}{2A}$$

-3.54138, 2.54138

Eigenvalues



- Conjecture 5: *There is one eigenvalue that is the roots of $x^2 + (-\mu)(c - 1)x - x^2$*

{9., -3.54138, -3.54138, 2.54138, 2.54138, -1., -1., -1., -1., -1., -1., -1., 0., 0., 0., 0., 0., 0.}

$$x^2 + (-\mu)(c - 1)x - x^2$$

$$1x^2 + (-1)(9 - 1)(1)x - 3^2$$

$$x^2 - 8x - 9$$

Irreducible: $(x^2 - 8x - 9)$

$$A = 1$$


$$B = -8$$

$$C = -9$$

$$\frac{(B \pm \sqrt{B^2 - 4AC})}{2A}$$

$$9, -1$$

Data



Thank you! Questions?